



二重装着を医療現場のスタンダードに！

DOUBLE GLOVING

血液や体液への接触を、本当に安全といえますか？

手術室で推奨されているパーソナルプロテクションエクイップメント（PPE）を装着していますか？



医療従事者の感染リスク低減のために、下記のガイドラインで二重装着を推奨しています



国立大学付属病院感染対策協議会病院感染対策ガイドライン（改訂第4版）



手術医療の実践ガイドライン改訂版（日本手術医学会）



RAPID ADVICE Guideline October 2014 issued by WHO（World Health Organization：世界保健機関）

(1) Murphy CL. The serious and ongoing issue of needlestick in Australian healthcare settings. Collegian 2013
(2) Partecke LI, Goerdts A-M, Langner I, et al. Incidence of microperforation for surgical gloves depends on duration of wear. Infect Control Hosp Epidemiol. 2009; 30:409-414
(3) Thomas S, Agarwal M, Mehta G. Intraoperative glove perforation—single versus double gloving in protection against skin contamination. Postgrad Med J. 2001; 77:458-460.
(4) Mischke C, Verbeek JH, Saarto A, Lavoie MC, Pahwa M, Ijaz S. Gloves, extra gloves special types of gloves for preventing percutaneous exposure injuries in healthcare personnel (Review) The Cochrane Collaboration 2014
(5) Berguer R, MD. Preventing Sharps Injuries in the Operating Room. Jour Am Coll Surg. 2004; 04:462-467.
(6) Laine T, Aarnio P. How often does glove perforation occur in surgery? Comparison between single gloves and a double-gloving system. American journal of surgery 2001; 181:564-6.
(7) Fry DE, Harris WE, Kohnke EN, Twomey CL. Influence of Double-Gloving on Manual Dexterity and Tactile Sensation of Surgeons. Journal of the American College of Surgeons. 3// 2010; 210(3):325-330.